

The Symplectic Geometry of Polygons in the 3-Sphere

Thomas Treloar

Abstract. We study the symplectic geometry of the moduli spaces $M_r = M_r(\mathbb{S}^3)$ of closed n -gons with fixed side-lengths in the 3-sphere. We prove that these moduli spaces have symplectic structures obtained by reduction of the fusion product of n conjugacy classes in $SU(2)$ by the diagonal conjugation action of $SU(2)$. Here the fusion product of n conjugacy classes is a Hamiltonian quasi-Poisson $SU(2)$ -manifold in the sense of [AKSM]. An integrable Hamiltonian system is constructed on M_r in which the Hamiltonian flows are given by bending polygons along a maximal collection of nonintersecting diagonals. Finally, we show the symplectic structure on M_r relates to the symplectic structure obtained from gauge-theoretic description of M_r . The results of this paper are analogues for the 3-sphere of results obtained for $M_r(\mathbb{H}^3)$, the moduli space of n -gons with fixed side-lengths in hyperbolic 3-space [KMT], and for $M_r(\mathbb{E}^3)$, the moduli space of n -gons with fixed side-lengths in \mathbb{E}^3 [KM1].

1 Introduction

This paper is an analogue to [KM1] and [KMT] which studied the symplectic geometry of moduli spaces of polygonal linkages with fixed side-lengths in Euclidean 3-space and hyperbolic 3-space, respectively. We obtain the moduli space of polygonal linkages with fixed side-lengths in the 3-sphere, \mathbb{S}^3 , by the reduction of a nondegenerate Hamiltonian quasi-Poisson $SU(2)$ -manifold in the sense of [AKSM].

We will use the following definitions throughout this paper. An (open) n -gon P in \mathbb{S}^3 is an ordered $(n + 1)$ -tuple of points in $\mathbb{S}^3 \subset \mathbb{C}^2$, $P = [y_1, \dots, y_{n+1}]$, called the *vertices*. We join the vertex y_i to the vertex y_{i+1} by a shortest geodesic segment e_i , called the *i -th edge*. This puts the restriction on the length of $e_i \leq \pi$ for all $1 \leq i \leq n$. Let $\text{Pol}_n(\mathbb{S}^3)$ denote the space of n -gons in \mathbb{S}^3 .

An n -gon is said to be *closed* if $y_{n+1} = y_1$. We let $\text{CPol}_n(\mathbb{S}^3)$ denote the space of closed n -gons in \mathbb{S}^3 . Let $\text{Isom}_+(\mathbb{S}^3)$ denote the group of orientation preserving isometries of \mathbb{S}^3 . There exists a natural (diagonal) action of $\text{Isom}_+(\mathbb{S}^3)$ on $\text{Pol}_n(\mathbb{S}^3)$ by

$$g \cdot [y_1, \dots, y_n] = [g \cdot y_1, \dots, g \cdot y_{n+1}].$$

Two n -gons $P = [y_1, \dots, y_{n+1}]$ and $P' = [y'_1, \dots, y'_{n+1}]$ are said to be *equivalent* if there exists $g \in \text{Isom}_+(\mathbb{S}^3)$ such that $g \cdot P = P'$.

Fix an n -tuple of strictly positive real numbers $r = (r_1, \dots, r_n) \in \mathbb{R}_+^n$. Denote the space of open n -gons with fixed side-lengths, $d(y_i, y_{i+1}) = r_i$, by $N_r(\mathbb{S}^3)$. Let $\text{CN}_r(\mathbb{S}^3) = N_r(\mathbb{S}^3) \cap \text{CPol}_n(\mathbb{S}^3)$, the space of closed polygons with fixed side-lengths. Finally, let $M_r(\mathbb{S}^3) = \text{CN}_r(\mathbb{S}^3) / \text{Isom}_+(\mathbb{S}^3)$. We study $M_r(\mathbb{S}^3)$, the moduli space

Received by the editors September 1, 2000; revised June 7, 2001.

Research partially supported by NSF grant DMS-98-03518.

AMS subject classification: 53D.

©Canadian Mathematical Society 2002.

closed n -gons with fixed side-lengths in \mathbb{S}^3 by the group of orientation preserving isometries.

Fix $* \in \mathbb{S}^3$. Denote by $\text{Rot}(\mathbb{S}^3, *) \subset \text{Isom}_+(\mathbb{S}^3)$ the group of rotations of \mathbb{S}^3 fixing $*$. Let $\text{Pol}_n(\mathbb{S}^3, *)$ denote the space of n -gons in \mathbb{S}^3 such that $y_1 = *$. Let $\text{CPol}_n(\mathbb{S}^3, *) = \text{CPol}_n(\mathbb{S}^3) \cap \text{Pol}_n(\mathbb{S}^3, *)$, $N_r(\mathbb{S}^3, *) = \text{Pol}_n(\mathbb{S}^3, *) \cap N_r(\mathbb{S}^3)$, and $\text{CN}_r(\mathbb{S}^3, *) = \text{CPol}_n(\mathbb{S}^3) \cap N_r(\mathbb{S}^3, *)$.

It is easy to see the space $M_r(\mathbb{S}^3)$ can be identified with $\text{CN}_r(\mathbb{S}^3, *) / \text{Rot}(\mathbb{S}^3, *)$, the moduli space of closed, based n -gons with fixed side-lengths in \mathbb{S}^3 by the group of rotations about the first vertex $*$.

The group of orientations preserving isometries is given by $\text{Isom}_+(\mathbb{S}^3) = (\text{SU}(2) \times \text{SU}(2)) / \{\pm I\}$. The group of rotations fixing the north and south poles is the diagonal subgroup, $K \simeq \text{PSU}(2)$, and translations are given by $\text{Isom}_+(\mathbb{S}^3) / K$ which we identify with $\text{SU}(2)$.

In this paper, a symplectic structure is obtained on $M_r(\mathbb{S}^3)$ by reduction of a Hamiltonian quasi-Poisson $\text{SU}(2)$ -manifold. We are interested in finding an integrable system on $M_r(\mathbb{S}^3)$. Denote by d_{ij} a shortest geodesic segment connecting the vertices y_i and y_j (we assume $i < j$), which we call a diagonal. Let ℓ_{ij} be the length of the diagonal d_{ij} . Then ℓ_{ij} is a continuous function on $M_r(\mathbb{S}^3)$, but it is not smooth when either $\ell_{ij} = 0$ or $\ell_{ij} = \pi$. If d_{ij} and d_{km} are nonintersecting diagonals, then

$$\{\ell_{ij}, \ell_{km}\} = 0.$$

By considering a maximal collection of nonintersecting diagonals, we obtain $\frac{1}{2} \dim(M_r(\mathbb{S}^3)) = 2(n-3)$ Poisson commuting Hamiltonians.

The Hamiltonian flow Ψ_{ij}^t associated to a ℓ_{ij} has the following nice description. Separate the polygon into two pieces via the diagonal d_{ij} . The Hamiltonian flow is given by leaving one piece fixed while rotating the other piece about the diagonal at constant angular velocity with period 2π . The flow Ψ_{ij}^t is called the *bending flow* along the diagonal d_{ij} and defines a \mathbb{T}^{n-3} -action on $M_r(\mathbb{S}^3)$.

The paper is organized as follows:

In Section 2, we give the Lie group description of spherical polygons.

In Section 3, we give criteria for the moduli space $M_r(\mathbb{S}^3)$ to be smooth and nonempty.

In Section 4, we give the necessary background material on quasi-Poisson manifolds.

In Section 5, we obtain a symplectic structure on $M_r(\mathbb{S}^3)$ and study the Hamiltonians ℓ_{ij} and their associated Hamiltonian flows.

In Section 6, we obtain the an action of the pure braid group on $M_r(\mathbb{S}^3)$ given by the time 1 Hamiltonian flows of a certain family of functions.

In Section 7, we relate the symplectic form on $M_r(\mathbb{S}^3)$ to symplectic form given on the relative character varieties on n -punctured 2-spheres.

We note that the moduli spaces of polygons in the 3-spaces of constant curvature give examples of completely integrable systems obtained from the theory of Manin pairs associated to a compact simple Lie group [AMM2]. The Manin pairs corresponding to the various moduli spaces are:

- $(\mathfrak{su}(2) \ltimes \mathfrak{su}(2)^*, \mathfrak{su}(2))$ for polygons in the zero curvature space (Lie-Poisson theory);
- $(\mathfrak{sl}_2(\mathbb{C}) = \mathfrak{su}(2)^{\mathbb{C}}, \mathfrak{su}(2))$ for polygons in negative curvature space (Poisson-Lie theory);
- $(\mathfrak{su}(2) \oplus \mathfrak{su}(2), \mathfrak{su}(2))$ for polygons in positive curvature space (quasi-Poisson Lie theory).

Acknowledgments The author would like to thank John Millson for introducing him to symplectic geometry and for many interesting discussions. Thanks are due to Bill Goldman for many useful conversations. The author would also like to thank Eckhard Meinrenken bringing to his attention [AKS] and [AKSM].

2 Lie Group Construction of Spherical Polygons

We identify $\mathbb{S}^3 \subset \mathbb{C}^2$ with $SU(2)$ by the map which takes (z_1, z_2) to $\begin{pmatrix} z_1 & -\bar{z}_2 \\ z_2 & \bar{z}_1 \end{pmatrix}$. Fix $* \in \mathbb{S}^3$ to be the north pole of \mathbb{S}^3 , $* = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. The group of orientation preserving isometries of \mathbb{S}^3 , $\text{Isom}_+(\mathbb{S}^3)$, is given by $G = (SU(2) \times SU(2)) / \{\pm I\}$ with the action of G on \mathbb{S}^3 given by

$$G \times \mathbb{S}^3 \rightarrow \mathbb{S}^3$$

$$((k_1, k_2), x) \mapsto k_1 x k_2^{-1}.$$

The diagonal subgroup, $K \simeq PSU(2)$, of G acts as the group of rotations on \mathbb{S}^3 fixing the north and south poles. Translations are then given by G/K , which we identify with $SU(2)$ by the map

$$G/K \rightarrow SU(2)$$

$$(k_1, k_2) \mapsto k_1 k_2^{-1}.$$

Recall the definitions of the various polygon spaces given in Section 1. We have a diffeomorphism from n copies of $SU(2)$ to $\text{Pol}_n(\mathbb{S}^3, *)$, the space of n -gons in \mathbb{S}^3 based at the point $*$ given by

$$\Phi: SU(2)^n \rightarrow \text{Pol}_n(\mathbb{S}^3, *)$$

$$(k_1, k_2, \dots, k_n) \mapsto [*, k_1*, k_1 k_2*, \dots, k_1 k_2 \cdots k_n*].$$

The condition for a polygon to be closed is $k_1 k_2 \cdots k_n = I$. The map Φ restricts to a diffeomorphism

$$\Phi: \{(k_1, \dots, k_n) \in SU(2)^n : k_1 \cdots k_n = I\} \rightarrow \text{CPol}_n(\mathbb{S}^3, *).$$

It is easily seen that the map Φ is K -equivariant where the action on $SU(2)^n$ is given by diagonal conjugation and on $\text{Pol}_n(\mathbb{S}^3, *)$ by the natural (diagonal) action.

We next see that fixing side-lengths for a polygon corresponds to restricting to conjugacy classes, $\mathcal{C}_\lambda \subset G$. Let $k, k' \in \mathcal{C}_\lambda$. If there exists $g \in K$ so that $g \cdot k = k'$, then

$$d(k \cdot *, *) = d(g \cdot k*, g \cdot *) = d(k' \cdot *, *)$$

We have the following lemma.

Lemma 2.1 *The map Φ induces a K -equivariant diffeomorphism between $\prod_{i=1}^n \mathcal{C}_{\lambda_i}$ and $N_r(\mathbb{S}^3, *)$, the configuration space of open based n -gon linkages with fixed side-lengths $r = (r_1, r_2, \dots, r_n) \in \mathbb{R}_+^n$, where $r_i = d(k_i*, *)$ for $k_i \in \mathcal{C}_{\lambda_i}$, $1 \leq i \leq n$.*

We now have the identification $\{(k_1, \dots, k_n) \in \prod_{i=1}^n \mathcal{C}_{\lambda_i} : k_1 \cdots k_n = I\} / \text{SU}(2)$ with $M_r(\mathbb{S}^3)$ by Φ .

3 Criteria For Smoothness and Nonemptiness

In this section we give necessary and sufficient conditions for the moduli space $M_r(\mathbb{S}^3)$ to be nonempty and sufficient conditions for $M_r(\mathbb{S}^3)$ to be a smooth manifold.

Let $\Pi: \text{CPol}_n(\mathbb{S}^3) \rightarrow \mathbb{R}_+^n$ be the map that assigns to an n -gon P its set of side-lengths, $\Pi(P) = (r_1, \dots, r_n)$ with $r_i = d(x_i, x_{i+1})$, $1 \leq i \leq n$. Let $I \subset \{1, 2, \dots, n\}$, \bar{I} denote the complement of I , $|I|$ denote the cardinality of I , and $r_I = \sum_{i \in I} r_i$.

Lemma 3.1 *The image of Π is the closed polyhedron D_n defined by the inequalities*

$$0 \leq r_i \leq \pi, \quad 1 \leq i \leq n, \quad \text{and} \\ r_I \leq r_{\bar{I}} + (|I| - 1)\pi, \quad I \subset \{1, 2, \dots, n\}, \quad \text{with } |I| \text{ odd.}$$

Proof The proof for n -gons in \mathbb{S}^2 was given by Galitzer [Ga]. Since any n -gon, P , in \mathbb{S}^m can be obtained from a finite number of bends along diagonals of an n -gon, P' , in $\mathbb{S}^2 \subset \mathbb{S}^n$, these inequalities hold for all $n \geq 2$. ■

We next give sufficient conditions for $M_r(\mathbb{S}^3)$ to be a smooth manifold. We will use two results and the notation from Section 5.1 (the reader will check that no circular reasoning is involved here). By Theorem 5.2 we find that $M_r(\mathbb{S}^3)$ is a symplectic manifold obtained by the reduction of a non-degenerate Hamiltonian quasi-Poisson manifold,

$$M_r(\mathbb{S}^3) \cong (\mu|_{N_r(\mathbb{S}^3, *)})^{-1}(1) / \text{SU}(2).$$

By Lemma 5.1, 1 is a regular value of μ unless there exists $P \in \text{CN}_r(\mathbb{S}^3, *)$ such that the infinitesimal isotropy $\text{su}_2|_P = \{X \in \text{su}_2 : X_{\text{CN}_r(\mathbb{S}^3, *)}(P) = 0\}$ is nonzero.

Definition 3.2 *An n -gon P is degenerate if it is contained in a geodesic.*

We now have the following lemma due to Galitzer [Ga], also see [KM3].

Lemma 3.3 $M_r(\mathbb{S}^3)$ is singular only if there exists a partition $\{1, \dots, n\} = I \amalg J$ with $\#(I) > 1, \#(J) > 1$ and $m \in \mathbb{Z}$ such that

$$\sum_{i \in I} r_i = \sum_{j \in J} r_j + 2m\pi.$$

Proof Clearly $\text{su}_2|_P = 0$ unless P is degenerate. But if P is degenerate there exists a partition $\{1, \dots, n\} = I \amalg J$ as above (I corresponds to the back-tracks and J to the forward-tracks of P). ■

Remark 3.4 In the terminology of [KM1], $M_r(\mathbb{S}^3)$ is smooth unless r is on a wall of D_n .

4 Quasi-Poisson Manifolds

We have seen that $N_r(\mathbb{S}^3, *)$ can be identified with the product of n conjugacy classes, $\prod_{i=1}^n \mathcal{C}_{\lambda_i}$. In this section, we introduce the machinery needed to construct quasi-Poisson bivectors on these spaces. We begin by reviewing basic definitions and results for quasi-Poisson K -spaces. For a complete treatment of quasi-Poisson manifolds see [AKS] and [AKSM].

4.1 Basic Definitions and the Moment Map

Let K be a Lie group whose Lie algebra \mathfrak{k} is equipped with an invariant nondegenerate bilinear form. Let $\{e_i\}$ be an orthonormal basis with respect to the bilinear form on \mathfrak{k} . We define $\varphi \in \wedge^3 \mathfrak{g}$ by

$$\varphi = \sum f_{ij}^k e_i \wedge e_j \wedge e_k,$$

where $[e_i, e_j] = \sum_k f_{ij}^k e_k$.

We denote by the subscript M, x_M , the vector field (resp. multivector field) on M induced by the action of K on M and $x \in \mathfrak{k}$ (resp. $x \in \wedge^j \mathfrak{k}$) satisfying

$$(1) \quad (x_M f)(m) = \left. \frac{d}{dt} \right|_{t=0} f(\exp(-tx) \cdot m)$$

where $f \in C^\infty(M)$ and $m \in M$. This is a Lie algebra homomorphism, i.e. $[x_M, y_M] = [x, y]_M$ for $x, y \in \mathfrak{k}$.

Definition 4.1 A quasi-Poisson manifold is a K -manifold M , equipped with an invariant bivector field $\pi_M \in C^\infty(M, \wedge^2 TM)$ such that the Schouten bracket of π_M is the invariant trivector field,

$$[\pi_M, \pi_M] = \varphi_M.$$

We next define the notion of a K -valued moment map.

Definition 4.2 An Ad-invariant map $\mu: M \rightarrow K$ is called a *moment map* for a quasi-Poisson K -manifold (M, π_M) if

$$\pi_M^\sharp(d(\mu^* f)) = (\mu^* (\frac{1}{2}(e^\lambda + e^\rho) f))_M,$$

for all functions $f \in C^\infty(K)$. The triple (M, π_M, μ) is then called a *Hamiltonian quasi-Poisson K -manifold*.

The following lemma gives us the formulation of the moment map most useful for this paper.

Lemma 4.3 Let (M, π_M) be a quasi-Poisson K -manifold. An Ad-equivariant map $\mu: M \rightarrow K$ is a moment map if and only if

$$\pi_M^\sharp(\mu^* \langle \theta, X \rangle) = \frac{1}{2}((1 + \text{Ad}_{\mu^{-1}})X)_M,$$

for all $X \in \mathfrak{k}$ and $\theta = k^{-1}dk$ the left-invariant Maurer-Cartan form on K .

Proof See [AKSM]. ■

Although the Schouten bracket of the bivector field on a Hamiltonian quasi-Poisson manifold is in general a nonzero invariant trivector field, we may still define a notion of reduction to obtain a symplectic manifold.

Lemma 4.4 (quasi-Poisson reduction) Let (M, π_M, μ) be a non-degenerate Hamiltonian quasi-Poisson manifold. Let M_* be the subset of M on which the K -action is free. Let $1 \in K$ be a regular value for $\mu: M \rightarrow K$. Then intersection of $\mu^{-1}(1)/K$ with M_*/K is a symplectic submanifold.

Proof See [AKSM]. ■

4.2 Conjugacy Classes as Quasi-Poisson Manifolds

The basic example of a Hamiltonian quasi-Poisson K -manifold is (K, π_K, μ) where the action is given by conjugation, the moment map $\mu = id_K$ is the identity map on K , and the bivector π_K is given by

$$\pi_K(k) = \frac{1}{2} \sum_i dR_k e_i \wedge dL_k e_i.$$

The bivector π_K restricts to a nondegenerate quasi-Poisson bivector on conjugacy classes $\mathcal{C} \subset K$. The triple $(\mathcal{C}, \pi_K|_{\mathcal{C}}, \mu|_{\mathcal{C}})$ is a Hamiltonian quasi-Poisson K -manifold.

4.3 Fusion Product of Quasi-Poisson Manifolds

Given Hamiltonian quasi-Poisson K -manifolds (M_1, π_1, μ_1) and (M_2, π_2, μ_2) , it is not true that $M_1 \times M_2$ with the product bivector is a Hamiltonian quasi-Poisson K -space for the diagonal action of K on $M_1 \times M_2$. We must construct a new bivector, π_{fus} , on $M_1 \times M_2$ for the diagonal action to be a quasi-Poisson action. $M_1 \times M_2$ with this bivector is called the *fusion product* and denoted $M_1 \circledast M_2$. This construction is due to [AKSM].

As defined in Section 4.1, the subscript M denotes the vector field, or multivector field, induced by the action of K on M . We define $\psi \in \wedge(\mathfrak{k} \oplus \mathfrak{k})$ to be

$$\psi = \frac{1}{2} \sum_i e_i^1 \wedge e_i^2$$

where $\{e_i\}$ is a basis of \mathfrak{k} and the superscripts refer to the respective \mathfrak{k} -summand.

Proposition 4.5 *Let (M, π) be a quasi-Poisson $K \times K \times H$ -manifold. Then*

$$\pi_{\text{fus}} = \pi - \psi_M$$

defines a quasi-Poisson structure on M for the diagonal $K \times H$ -action. Moreover, if $(\mu_1, \mu_2, \mu_H): M \rightarrow K \times K \times H$ is a moment map for the action, then the point-wise product $(\mu_1 \mu_2, \mu_H)$ is a moment map for the diagonal $K \times H$ -action.

Proof See [AKSM, Proposition 5.1].

In the previous section, we showed a conjugacy class $\mathcal{C} \subset K$ was a Hamiltonian quasi-Poisson manifold. For this paper, the Hamiltonian quasi-Poisson spaces we are most interested in are the fusion products of n conjugacy classes in K .

Example 4.6 Let $(\mathcal{C}_{\lambda_1}, \pi_1, \mu_1)$ and $(\mathcal{C}_{\lambda_2}, \pi_2, \mu_2)$ be conjugacy classes in K . Then $\mathcal{C}_{\lambda_1} \times \mathcal{C}_{\lambda_2}$ with the bivector

$$\pi_{\text{fus}}(k_1, k_2) = \pi_1(k_1) + \pi_2(k_2) - \sum_i (dL_{k_1} e_i^1 - dR_{k_1} e_i^1) \wedge (dL_{k_2} e_i^2 - dR_{k_2} e_i^2),$$

where the superscripts denote the conjugacy class on which e_i acts, is a Hamiltonian quasi-Poisson K -space where the action is given by diagonal conjugation, $k \cdot (k_1, k_2) = (kk_1k^{-1}, kk_2k^{-1})$. The moment map associated to this action is the product $\mu(k_1, k_2) = k_1 k_2$.

We are now in the position to give a formula for quasi-Poisson bivector on the product of n conjugacy classes given by fusion. The fusion product $\circledast_{i=1}^n \mathcal{C}_{\lambda_i}$ is a Hamiltonian quasi-Poisson K -space with action given by the diagonal conjugation and moment map $\mu: \circledast_{i=1}^n \mathcal{C}_{\lambda_i} \rightarrow K$ given by the product, $\mu(k_1, k_2, \dots, k_n) = k_1 k_2 \cdots k_n$. The quasi-Poisson bivector on this space is given by

$$\pi_{\text{fus}} = \frac{1}{2} \sum_i \sum_l (dR_{k_i} e_l^i \wedge dL_{k_i} e_l^i) - \frac{1}{2} \sum_{i < j} \sum_l (dL_{k_i} e_l^i - dR_{k_i} e_l^i) \wedge (dL_{k_j} e_l^j - dR_{k_j} e_l^j)$$

where the superscripts i, j denote the conjugacy class $\mathcal{C}_{\lambda_i}, \mathcal{C}_{\lambda_j} \subset \otimes_{i=1}^n \mathcal{C}_{\lambda_i}$ on which $e_l \in \mathfrak{k}$ acts.

The following remark from [AKS, Example 5.5.4] gives the nondegeneracy of $\otimes_{i=1}^n \mathcal{C}_{\lambda_i}$.

Remark 4.7 Let (M, π, μ) be a Hamiltonian quasi-Poisson K space. Then (M, π, μ) is nondegenerate if and only if for each $m \in M$,

$$\ker(\pi^\sharp(m)) = \{\mu^*(x, \theta) : x \in \ker(1 + \text{Ad}_{\mu(m)})\}$$

where $x \in \mathfrak{k}$.

4.4 Poisson Bracket on $C^\infty(\otimes_{i=1}^n \mathcal{C}_{\lambda_i})^K$

For a general quasi-Poisson manifold (M, π_M) , we can define a bracket on $C^\infty(M)$ by $\{\psi_1, \psi_2\} = \pi_M(d\psi_1, d\psi_2)$, where $\psi_1, \psi_2 \in C^\infty(M)$. This bracket is not, in general, a Poisson bracket. This is because the Schouten bracket of π_M is an invariant trivector field, φ_M , not necessarily zero. The bracket does however define a Poisson bracket when we restrict to $C^\infty(M)^K$, the smooth K -invariant functions on M .

We will next find a formula for the Poisson bracket on the K -invariant functions on the fusion product of n conjugacy classes in K . Let $\mathfrak{k}_i \subset \bigoplus_{i=1}^n \mathfrak{k}$ denote the i -th summand, $\nu_i \in \mathfrak{k}_i$, and $k = (k_1, \dots, k_n) \in \otimes_{i=1}^n \mathcal{C}_{\lambda_i}$. For $\psi \in C^\infty(\otimes_{i=1}^n \mathcal{C}_{\lambda_i})$ we define

$$D_i \psi: \otimes_{i=1}^n \mathcal{C}_{\lambda_i} \rightarrow \mathfrak{k}_i, \quad D'_i \psi: \otimes_{i=1}^n \mathcal{C}_{\lambda_i} \rightarrow \mathfrak{k}_i$$

by

$$\begin{aligned} \langle D_i \psi(k), \nu_i \rangle &= \left. \frac{d}{dt} \right|_{t=0} \psi(k_1, \dots, e^{t\nu_i} k_i, \dots, k_n) \\ \langle D'_i \psi(k), \nu_i \rangle &= \left. \frac{d}{dt} \right|_{t=0} \psi(k_1, \dots, k_i e^{t\nu_i}, \dots, k_n). \end{aligned}$$

Here $\langle \cdot, \cdot \rangle$ is the nondegenerate invariant bilinear form on \mathfrak{k} extended to $\bigoplus_{i=1}^n \mathfrak{k}$ by $\langle x, y \rangle = \sum_{i=1}^n \langle x_i, y_i \rangle$ for $x = (x_1, \dots, x_n) \in \bigoplus_{i=1}^n \mathfrak{k}$ and $y = (y_1, \dots, y_n) \in \bigoplus_{i=1}^n \mathfrak{k}$.

Lemma 4.8

$$\text{Ad}_{k_i} D'_i \psi(k) = D_i \psi(k).$$

Proof

$$\begin{aligned} \langle \text{Ad}_{k_i} D'_i \psi(k), \nu_i \rangle &= \langle D'_i \psi(k), \text{Ad}_{k_i^{-1}} \nu_i \rangle \\ &= \left. \frac{d}{dt} \right|_{t=0} \psi(k_1, \dots, k_i \text{Ad}_{k_i^{-1}} e^{t\nu_i}, \dots, k_n) \\ &= \left. \frac{d}{dt} \right|_{t=0} \psi(k_1, \dots, e^{t\nu_i} k_i, \dots, k_n) \\ &= \langle D_i \psi(k), \nu_i \rangle. \quad \blacksquare \end{aligned}$$

We also define

$$\Psi_j(k) = \sum_{i=1}^{j-1} [D_i\psi(k) - D'_i\psi(k)] + D_j\psi(k)$$

We now give a formula for the Poisson bracket on $C^\infty(\otimes_{i=1}^n \mathcal{C}_{\lambda_i})^K$.

Proposition 4.9 *Let $\phi, \psi \in C^\infty(\otimes_{i=1}^n \mathcal{C}_{\lambda_i})^K$ then*

$$\{\phi, \psi\}(k) = \sum_{j=1}^n \langle D'_j\phi(k) - D_j\phi(k), \Psi_j(k) \rangle.$$

Proof Let us first note that $\sum_l \langle x, e_l \rangle \langle y, e_l \rangle = -\langle x, y \rangle$ for $x, y \in \mathfrak{k}$ and $\{e_l\}$ an orthonormal basis of \mathfrak{k} . Now,

$$\begin{aligned} \{\phi, \psi\}(k) &= \pi_{\text{fus}}(d\phi, d\psi) \\ &= \frac{1}{2} \sum_i \sum_l (dR_k e_l^i \wedge dL_k e_l^i)(d\phi, d\psi) \\ &\quad - \frac{1}{2} \sum_{i < j} \sum_l ((dL_k e_l^i - dR_k e_l^i) \wedge (dL_k e_l^j - dR_k e_l^j))(d\phi, d\psi) \\ &= \frac{1}{2} \sum_i \sum_l \langle D_i\phi, e_l^i \rangle \langle D'_i\psi, e_l^i \rangle - \langle D'_i\phi, e_l^i \rangle \langle D_i\psi, e_l^i \rangle \\ &\quad - \frac{1}{2} \sum_{i < j} \sum_l \langle D'_i\phi - D_i\phi, e_l^i \rangle \langle D'_j\psi - D_j\psi, e_l^j \rangle \\ &\quad + \frac{1}{2} \sum_{i < j} \sum_l \langle D'_j\phi - D_j\phi, e_l^j \rangle \langle D'_i\psi - D_i\psi, e_l^i \rangle \\ &= \frac{1}{2} \sum_i \langle D'_i\phi, D_i\psi \rangle - \langle D_i\phi, D'_i\psi \rangle \\ &\quad + \frac{1}{2} \sum_{i < j} \langle D'_i\phi - D_i\phi, D'_j\psi - D_j\psi \rangle - \langle D'_j\phi - D_j\phi, D'_i\psi - D_i\psi \rangle \\ &= \frac{1}{2} \sum_i \langle D'_i\phi, D_i\psi \rangle - \langle D_i\phi, D'_i\psi \rangle \\ &\quad + \frac{1}{2} \sum_{i < j} \langle D'_i\phi + D_i\phi, D'_j\psi - D_j\psi \rangle - \sum_{i > j} \langle D'_i\phi - D_i\phi, D'_j\psi - D_j\psi \rangle. \end{aligned}$$

But since $\psi \in C^\infty(\otimes_{i=1}^n \mathcal{C}_{\lambda_i})^K$ is K -invariant, a quick calculation shows

$$\sum_i [D_i\psi - D'_i\psi] = 0.$$

Using this fact and also that $\langle D'_i \phi, D'_i \psi \rangle = \langle D_i \phi, D_i \psi \rangle$ for all i , we can rewrite the above as,

$$\begin{aligned} \{\phi, \psi\} &= \frac{1}{2} \sum_i \langle D'_i \phi - D_i \phi, D_i \psi + D'_i \psi \rangle \\ &\quad - \frac{1}{2} \sum_{i \geq j} \langle D'_i \phi - D_i \phi, D'_j \psi - D_j \psi \rangle - \frac{1}{2} \sum_{i > j} \langle D'_i \phi - D_i \phi, D'_j \psi - D_j \psi \rangle \\ &= \sum_i \langle D'_i \phi - D_i \phi, \Psi_i \rangle. \quad \blacksquare \end{aligned}$$

From the above proposition we can also define the Hamiltonian vector field X_ψ associated to $\psi \in C^\infty(\otimes_{i=1}^n \mathcal{C}_{\lambda_i})^K$ by $X_\psi = \pi^\sharp(d\psi)$.

Corollary 4.10 *The Hamiltonian vector field $X_\psi(k) = (X_1(k), \dots, X_n(k))$ associated to the K -invariant function $\psi \in C^\infty(\otimes_{i=1}^n \mathcal{C}_{\lambda_i})^K$ is given by*

$$X_j(k) = dL_{k_j} \Psi_j - dR_{k_j} \Psi_j, \quad 1 \leq j \leq n.$$

Proof We use the convention $\{\phi, \psi\} = d\phi(X_\psi) = \sum_{j=1}^n d_j \phi(X_j(k))$. Proposition 4.9 gives us

$$\begin{aligned} d\phi(X_\psi(k)) &= \{\phi, \psi\} \\ &= \sum_{j=1}^n \langle D'_j \phi - D_j \phi, \Psi_j \rangle \\ &= \sum_{j=1}^n d_j \phi(dL_{k_j} \Psi_j) - d_j \phi(dR_{k_j} \Psi_j) \\ &= \sum_{j=1}^n d_j \phi(dL_{k_j} \Psi_j - dR_{k_j} \Psi_j). \quad \blacksquare \end{aligned}$$

5 Integrable Systems on $M_r(\mathbb{S}^3)$

We restrict to the case which gives rise to $M_r(\mathbb{S}^3)$, that is $K = \text{SU}(2)$ and $\langle \cdot, \cdot \rangle = -\frac{1}{2} \text{Trace}(\cdot)$, although most of the results of this section follow for $K = \text{SU}(n)$ and $\langle \cdot, \cdot \rangle$ the Killing form.

5.1 $M_r(\mathbb{S}^3)$ as a Symplectic Manifold

In Section 2, we constructed the K -equivariant diffeomorphism $\Phi: \prod_{i=1}^n \mathcal{C}_{\lambda_i} \rightarrow N_r(\mathbb{S}^3, *)$, defined by

$$\Phi(k_1, \dots, k_n) = [*, k_1*, k_1 k_2*, \dots, k_1 k_2 \cdots k_n*].$$

In Section 4.4, we constructed a nondegenerate quasi-Poisson structure on $\prod_{i=1}^n \mathcal{C}_{\lambda_i}$ with moment map associated to diagonal conjugation given by $\mu(k_1, k_2, \dots, k_n) = k_1 k_2 \cdots k_n$. By Lemma 4.4, if 1 is a regular value of μ we obtain a symplectic structure on $\mu^{-1}(1)/K$ given by reduction.

Lemma 5.1 *1 is a regular value of μ if and only if $\{x \in \mathfrak{k} : x_{\prod_{i=1}^n \mathcal{C}_{\lambda_i}} = 0\} = 0$ for all $k \in \mu^{-1}(1)$.*

Proof We refer to Lemma 4.3. Let $x \in \mathfrak{k}$ and $k \in \prod_{i=1}^n \mathcal{C}_{\lambda_i}$. Then

$$\begin{aligned} x \in (\operatorname{Im}(d\mu|_k))^\perp &\Leftrightarrow \mu^* \langle x, \theta \rangle = 0 \\ &\Leftrightarrow 0 = \pi^\sharp(\mu^* \langle x, \theta \rangle) = x_{\prod_{i=1}^n \mathcal{C}_{\lambda_i}}. \quad \blacksquare \end{aligned}$$

By Lemma 3.3, a polygon is said to be *degenerate* if it can be contained in a geodesic of \mathbb{S}^3 . It follows from the above lemma that if there does not exist $k \in \mu^{-1}(1) \subset \prod_{i=1}^n \mathcal{C}_{\lambda_i}$ such that $\Phi(k)$ is a degenerate polygon, then 1 is a regular value of μ .

We can therefore construct a symplectic structure on $\mu^{-1}(1)/K$ by quasi-Poisson reduction.

Theorem 5.2 *The moduli space $M_r(\mathbb{S}^3)$ containing no degenerate polygons has a symplectic structure which is obtained from the symplectic structure on $\mu^{-1}(1)/K$ via the diffeomorphism Φ .*

In [AKSM], the authors prove the correspondence between nondegenerate quasi-Poisson K -manifolds and quasi-Hamiltonian K -manifolds in the sense of [AMM1]. In Section 7, we need a formula for the 2-form on $\prod_{i=1}^n \mathcal{C}_{\lambda_i}$ which corresponds to π_{fus} . We will relate this 2-form to the 2-form obtained from the gauge-theoretic description of $M_r(\mathbb{S}^3)$.

Remark 5.3 The 2-form on $\prod_{i=1}^n \mathcal{C}_{\lambda_i}$ which corresponds to π_{fus} is given by

$$\tilde{\omega} = \sum_{i=1}^n \omega_i + \frac{1}{2} \sum_{i=1}^n \sum_{j=i+1}^n (\operatorname{Ad}_{k_1 \cdots k_{i-1}} \bar{\theta}_i \wedge_b \operatorname{Ad}_{k_1 \cdots k_{j-1}} \bar{\theta}_j).$$

where ω_i is the quasi-Hamiltonian 2-form on the conjugacy class $C_i \subset \operatorname{SU}(2)$, see [AMM1], and $\bar{\theta}_i$ is the right-invariant Maurer-Cartan form on $C_i \subset \operatorname{SU}(2)$. We denote by \wedge_b the wedge product together with the Killing form on K .

5.2 Hamiltonian Vector Fields

Let d_i denote the diagonal connecting the 1-st vertex with the $(i+1)$ -th vertex. Let ℓ_i be the function giving the length of d_i . We show that $\{\ell_i\}_{i=2}^{n-1}$ give us an integrable system on $M_r(\mathbb{S}^3)$. We first consider the functions

$$f_i(k) = \operatorname{tr}(k_1 \cdots k_i), \quad 1 \leq i \leq n.$$

They are related to ℓ_i by

$$\ell_i = \cos^{-1}\left(-\frac{1}{2}f_i\right).$$

In this section we compute the Hamiltonian vector fields X_{f_i} associated to the functions f_i .

See Section 4.4 for the definition of the Poisson bracket on $C^\infty(\otimes_{i=1}^n \mathcal{C}_{\lambda_i})^K$. We leave it to the reader to verify the following lemma.

Lemma 5.4

$$\begin{aligned} D_{i+1}f_j(k) &= D'_i f_j(k), \quad 1 \leq i \leq j-1 \\ D_1 f_j(k) &= D'_j f_j(k) \end{aligned}$$

for all $1 \leq j \leq n$.

We define $F_j: \otimes_{i=1}^n \mathcal{C}_{\lambda_i} \rightarrow \mathfrak{k}$ by

$$F_j(k) = ((k_1 \cdots k_j) - (k_1 \cdots k_j)^{-1}).$$

We then have the following lemma.

Lemma 5.5 $F_j(k) = -D_1 f_j(k)$

Proof For $k \in \otimes_{i=1}^n \mathcal{C}_{\lambda_i}$ and $X \in \mathfrak{k}$

$$\begin{aligned} \langle D_1 f_j(k), X \rangle &= \left. \frac{d}{dt} \right|_{t=0} \text{tr}(e^{tX} k_1 k_2 \cdots k_j) \\ &= \text{tr}(X k_1 k_2 \cdots k_j) \\ &= \text{tr}(k_1 k_2 \cdots k_j X) \end{aligned}$$

but since

$$\text{tr}((k_1 k_2 \cdots k_j)^{-1} X) = \text{tr}((k_1 \cdots k_j)^* X) = \text{tr}(X^* k_1 \cdots k_j) = -\text{tr}(k_1 \cdots k_j X)$$

it follows that

$$\begin{aligned} \text{tr}(k_1 k_2 \cdots k_j X) &= \frac{1}{2} \text{tr}\left(\left((k_1 k_2 \cdots k_j) - (k_1 \cdots k_j)^{-1}\right) X\right) \\ &= \langle -(k_1 \cdots k_j) + (k_1 \cdots k_j)^{-1}, X \rangle. \end{aligned}$$

Since $-(k_1 \cdots k_j) + (k_1 \cdots k_j)^{-1} \in \mathfrak{k}$ and $\langle \cdot, \cdot \rangle$ is a nondegenerate bilinear form, we have $D_1 f_j(k) = -((k_1 \cdots k_j) - (k_1 \cdots k_j)^{-1}) = -F_j(k)$. \blacksquare

We have the following formula of the Hamiltonian vector fields X_{f_i} .

Theorem 5.6 *The Hamiltonian vector field X_{f_i} has an i -th component given by*

$$\begin{aligned} (X_{f_j}(k))_i &= dR_{k_i}F_j(k) - dL_{k_i}F_j(k), \quad 1 \leq i \leq j, \\ (X_{f_j}(k))_i &= 0, \quad j < i \leq n \end{aligned}$$

Proof Recall from Corollary 4.10 that for $\psi \in C^\infty(\otimes_{i=1}^n \mathcal{C}_{\lambda_i})^K$, $X_\psi(k)$ is given by

$$(X_\psi(k))_i = dL_{k_i}\Psi_i(k) - dR_{k_i}\Psi_i(k)$$

where $\Psi_i(k) = D_1\psi(k) - D'_1\psi(k) + D_2\psi(k) - \dots - D_{i-1}\psi(k) + D_i\psi(k)$. This together with Lemma 5.4 gives us

$$(X_{f_j}(k))_i = dL_{k_i}D_1f_j(k) - dR_{k_i}D_1f_j(k), \quad 1 \leq i \leq j$$

and

$$(X_{f_j}(k))_i = 0, \quad j < i \leq n.$$

In Lemma 5.5 we obtained $-F_j(k) = D_1f_j(k)$ completing the proof. \blacksquare

5.3 Commuting Hamiltonians

In this section we will show the family of Hamiltonians under consideration, $\{f_j\}_{j=1}^n$, Poisson commute.

Proposition 5.7 $\{f_i, f_j\} \equiv 0$ for all i, j .

Proof Without loss of generality we may assume $i < j$, then by Proposition 4.9

$$\begin{aligned} \{f_i, f_j\}(k) &= \sum_{k=1}^j \langle D'_k f_i(k) - D_k f_i(k), F_j(k) \rangle \\ &= - \left\langle \sum_{k=1}^j (D'_k f_i(k) - D_k f_i(k)), F_j(k) \right\rangle \\ &= \langle 0, F_j(k) \rangle \\ &= 0 \end{aligned}$$

Here we use $\sum_{k=1}^i (D_k f_j - D'_k f_j) = 0$. \blacksquare

5.4 Hamiltonian Flow

In this section we will calculate the Hamiltonian flow, ϱ_j^t , associated to f_j . We will see that these flows are the bending flows described in the introduction. The Hamiltonian flow is the solution to the ODE

$$(2) \quad \begin{aligned} \frac{dk_i}{dt} &= dR_{k_i}F_j(k) - dL_{k_i}F_j(k) = [F_j(k), k_i], \quad 1 \leq i \leq j, \\ \frac{dk_i}{dt} &= 0, \quad j < i \leq n. \end{aligned}$$

Since we are working with matrix groups, we use the matrix commutator $[\cdot , \cdot]$ in the above equation.

Lemma 5.8 $F_j(k)$ is invariant along solution curves of (2).

Proof To prove the lemma, it suffices to show that $\tilde{\varrho}_j^t(k) = k_1(t) \cdots k_j(t)$ is invariant along solution curves of (2).

$$\begin{aligned} \frac{d}{dt} \tilde{\varrho}_j^t(k) &= \frac{d}{dt} (k_1(t)k_2(t) \cdots k_j(t)) \\ &= \frac{dk_1}{dt}(t)k_2(t) \cdots k_{j-1}(t)k_j(t) + \cdots + k_1(t)k_2(t) \cdots \frac{dk_j}{dt}(t) \\ &= [F_j(k(t)), k_1(t)] k_2(t) \cdots k_j(t) + k_1(t)k_2(t) \cdots [F_j(k(t)), k_j(t)] \\ &= F_j(k(t)) k_1(t) \cdots k_j(t) - k_1(t) \cdots k_j(t) F_j(k(t)) \\ &= 0. \end{aligned} \quad \blacksquare$$

Lemma 5.9 The curve $\exp(tF_j(k)) \subset K$ is periodic with period $2\pi/\sqrt{4-f_j^2}$.

Proof To simplify notation, let $X = F_j(k) \in \mathfrak{k}$. Then

$$X^{-1} = -\frac{1}{\det(X)}X$$

giving us

$$X^2 = -(\det(X))X^{-1}X = -\det(X)I.$$

So,

$$\begin{aligned} \exp tX &= \sum_{n=0}^{\infty} \frac{t^n X^n}{n!} \\ &= \sum_{n=1}^{\infty} \frac{(-1)^n (t \det(X))^n}{(2n)!} I + \sum_{n=1}^{\infty} \frac{(-1)^n (t \det(X))^n}{(2n+1)!} \frac{X}{\sqrt{\det(X)}} \\ &= \cos(t\sqrt{\det(X)}) I + \frac{\sin(t\sqrt{\det(X)})}{\sqrt{\det(X)}} X \\ &= \cos(t\sqrt{4-f_j(k)^2}) I + \frac{\sin(t\sqrt{4-f_j(k)^2})}{\sqrt{4-f_j(k)^2}} F_j(k). \end{aligned}$$

Therefore the curve is periodic with period $2\pi/\sqrt{4-f_j(b)^2}$. ■

We are now able to calculate the formula for the Hamiltonian flow ϱ_j^t .

Theorem 5.10 Suppose $P \in M_r(\mathbb{S}^3)$ has vertices given by $[*, k_1*, \dots, k_1 \cdots k_n*]$. Then the Hamiltonian flow, $P(t) = \varrho_j^t(P)$, associated to the Hamiltonian f_j has vertices given by $P(t) = [*, \tilde{k}_1(t)*, \dots, \tilde{k}_n(t)*]$ where

$$\tilde{k}_i(t) = \begin{cases} \text{Ad}_{\exp(tF_j(k))}(k_1 \cdots k_i), & 1 \leq i < j \\ k_1 \cdots k_i, & j \leq i \leq n. \end{cases}$$

The flow is periodic with period $2\pi/\sqrt{4-f_j^2}$.

The flow $\varrho_j^t(P)$ has the following geometric description. Let d_j be the diagonal connecting the first vertex with the $(j+1)$ -th vertex, that is $*$ with $k_1 \cdots k_j*$. Then $\varrho_j^t(P)$ rotates the first j vertices, $k_1 \cdots k_{i-1}*$, for $2 \leq i \leq j$, about the diagonal d_j at constant angular velocity. The flows $\{\varrho_j^t\}$, $1 < j < n$, do not give rise to a torus action on $M_r(\mathbb{S}^3)$ since they do not have constant period. For example, as the length of a diagonal goes to zero, the period of flow about that diagonal goes to infinity.

To get a torus action on $M_r(\mathbb{S}^3)$, we need to look instead at the length functions $\ell_j(k) = \cos^{-1}(-\frac{1}{2}f_j(k))$. Then

$$d\ell_j = \frac{1}{\sqrt{4-f_j^2}}df_j$$

and

$$X_{\ell_j} = \frac{1}{\sqrt{4-f_j^2}}X_{f_j}.$$

It is not difficult to see that the family of functions $\{\ell_j\}_{j=2}^{n-2}$ also Poisson commute, although the Hamiltonian flows for these functions are not everywhere defined on $M_r(\mathbb{S}^3)$. We restrict to the space $M'_r(\mathbb{S}^3)$ such $\ell_j \neq 0$ or $\ell_j \neq \pi$ for all $1 \leq j \leq n$. The Hamiltonian flows $\{\Psi_j^t\}$ on $M'_r(\mathbb{S}^3)$ associated to $\{\ell_j\}$ are periodic with constant period 2π and constant angular velocity. These flows define a Hamiltonian $(n-3)$ -torus action on the space $M'_r(\mathbb{S}^3)$.

6 Braid Action on $M_r(\mathbb{S}^3)$

There exists an action of the pure braid group \mathcal{P}_n on the manifold $M_r(\mathbb{S}^3)$ which preserves the symplectic structure. In this section, we show that the generators of the pure braid group arise as the time 1 Hamiltonian flows of the family of functions h_{ij} , $1 \leq i < j \leq n-1$ where $h_{ij} \in C^\infty(M_r(\mathbb{S}^3))^K$ is defined by,

$$h_{ij}(k) = \frac{1}{2} \left(\cos^{-1} \left(-\frac{1}{2} \text{tr}(k_i k_j) \right) \right)^2.$$

Let π_{12} denote the quasi-Poisson bivector on $\mathcal{C}_1 \otimes \mathcal{C}_2$. We have the following proposition.

Proposition 6.1 [AKSM, Proposition 5.7] *The diffeomorphism $R: C_1 \otimes C_2 \rightarrow C_2 \otimes C_1$ given by $R(k_1, k_2) = (\text{Ad}_{k_1} k_2, k_1)$ is a bivector map taking π_{12} to π_{21} .*

A similar proof gives us,

Proposition 6.2 *The diffeomorphism $R': C_1 \otimes C_2 \rightarrow C_2 \otimes C_1$ given by $R'(k_1, k_2) = (k_2, \text{Ad}_{k_2^{-1}} k_1)$ is also a bivector map taking π_{12} to π_{21} .*

Remark 6.3 $R \circ R' = \text{Id}_{C_1 \otimes C_2} = R' \circ R$

We now define $R_i: C_1 \otimes \cdots \otimes (C_i \otimes C_{i+1}) \otimes \cdots \otimes C_n \rightarrow C_1 \otimes \cdots \otimes (C_{i+1} \otimes C_i) \otimes \cdots \otimes C_n$ to be the map given by

$$R_i(k_1, \dots, k_i, k_{i+1}, \dots, k_n) = (k_1, \dots, \text{Ad}_{k_i} k_{i+1}, k_i, \dots, k_n)$$

that is, R applied to the i -th and $(i+1)$ -th term of $M_r(\mathbb{S}^3)$. R'_i can be defined in a similar way. See [Bi] for definitions of the full braid group, \mathcal{B}_n , and the pure braid group, \mathcal{P}_n .

Lemma 6.4 *The full braid group \mathcal{B}_n has a faithful representation as a group of automorphism of the closed n -gons in \mathbb{S}^3 in which side-lengths are fixed but the order of the sides is not fixed. The generators of \mathcal{B}_n are given by R_i , $1 \leq i \leq n-1$.*

We now restrict \mathcal{B}_n to \mathcal{P}_n to get an action of the pure braid group on $\otimes_{i=1}^n \mathcal{C}_{\lambda_i}$. This action induces a symplectomorphism on the moduli space $M_r(\mathbb{S}^3)$.

Corollary 6.5 *Let $A_{ij} = R_{j-1} \circ \cdots \circ R_{i+1} \circ R_i^2 \circ R'_{i+1} \circ \cdots \circ R'_{j-1}$, $1 \leq i < j \leq n$. A_{ij} induces a symplectomorphism from $M_r(\mathbb{S}^3)$ to itself. The A_{ij} , $1 \leq i < j \leq n$ are generators of \mathcal{P}_n which has a faithful representation as a group of automorphisms of $M_r(\mathbb{S}^3)$.*

We will now show that the braid group actions A_{ij} can be realized as the time one Hamiltonian flows of the Hamiltonians h_{ij} given at the beginning of this section. We first study the Hamiltonian flows associated to the functions $f_{ij} \in C^\infty(\otimes_{i=1}^n \mathcal{C}_{\lambda_i})^K$ given by $f_{ij}(k) = \text{tr}(k_i k_j)$. Define $F_{ij}: \otimes_{i=1}^n \mathcal{C}_{\lambda_i} \rightarrow \mathfrak{k}$ by $F_{ij}(k) = ((k_i k_j) - (k_i k_j)^{-1})$.

The Hamiltonian flow associated to f_{ij} is given by $\Phi_{ij}^t(k) = (\widehat{k}_1(t), \dots, \widehat{k}_n(t))$ where

$$\widehat{k}_l(t) = \begin{cases} k_l, & 0 < l < i \text{ and } j < l < n+1 \\ \text{Ad} \left(\exp(t F_{ij}(k)) \right) k_l, & l = i, j \\ \text{Ad} \left(\exp(t F_{ij}(k)) k_j \exp(-t F_{ij}(k)) k_j^{-1} \right) k_l, & i < l < j. \end{cases}$$

Following the proof of Lemma 5.9, we obtain

Lemma 6.6

$$\exp \left(\frac{\cos^{-1} \left(-\frac{1}{2} \text{tr}(k) \right)}{\sqrt{4 - \text{tr}^2(k)}} (k - k^{-1}) \right) = k$$

We now notice that for time $t = \frac{\cos^{-1}\left(-\frac{1}{2}f_{ij}(k)\right)}{\sqrt{4-f_{ij}^2(k)}}$,

$$\Phi_{ij}^t = A_{ij}.$$

The time for which Φ_{ij}^t flows depends on the point in $M_r(\mathbb{S}^3)$ at which flow begins. We would like this time to be independent of the starting point. This can be achieved by taking functions $h_{ij} = \frac{1}{2}\left(\cos^{-1}\left(-\frac{1}{2}f_{ij}\right)\right)^2$. The Hamiltonian flow $\tilde{\Phi}_{ij}^t$ associated to h_{ij} is the renormalization of the flow Φ_{ij}^t so that

$$\tilde{\Phi}_{ij}^1 = A_{ij}.$$

We can see the pure braid group as the integer points in the Hamiltonian flows $\tilde{\Phi}_{ij}^t$, $1 \leq i < j \leq n$.

7 Connection With Symplectic Forms on Relative Character Varieties of n -Punctured 2-Spheres

In this section, we relate the symplectic form on $M_r(\mathbb{S}^3)$ given in Remark 5.3 to the symplectic form of Goldman type obtained from the description of $M_r(\mathbb{S}^3)$ as the moduli space of flat connections on an n -punctured 2-sphere. We follow the arguments of Kapovich and Millson [KM1, Section 5] which considers the analogous question for $M_r(\mathbb{E}^3)$. As a consequence, we obtain, using a result of L. Jeffrey, a symplectomorphism from $M_r(\mathbb{E}^3)$ and $M_r(\mathbb{S}^3)$ for sufficiently small side-lengths.

We begin with the general case in which G is any Lie group with Lie algebra \mathfrak{g} which admits a nondegenerate, G -invariant, symmetric, bilinear form.

7.1 Relative Characteristic Varieties and Parabolic Cohomology

Let $\Sigma = \mathbb{S}^2 - \{p_1, \dots, p_n\}$ denote the n -punctured 2-sphere and U_1, \dots, U_n be disjoint open disc neighborhoods of p_1, \dots, p_n , respectively. Further, Γ is the fundamental group of Σ with generators γ_i and $T = \{\Gamma_1, \dots, \Gamma_n\}$ is the collection of subgroups of Γ with Γ_i the cyclic subgroup generated by γ_i .

Fix $\rho_0 \in \text{Hom}(\Gamma, G)$ a representation. In [KM1], the relative representation variety $\text{Hom}(\Gamma, T; G)$ is defined as the representations $\rho: \Gamma \rightarrow G$ such that $\rho|_{\Gamma_i}$ is contained in the closure of the conjugacy class of $\rho_0|_{\Gamma_i}$.

Remark 7.1 If $G = \text{SU}(2)$, there exists a ρ_0 such that the relative character variety $\text{Hom}(\Gamma, T; G)/G$ is isomorphic to $M_r(\mathbb{S}^3)$. We will make this isomorphism explicit later on.

Let $\rho \in \text{Hom}(\Gamma, T; G)$. Then ρ induces a flat principal G -bundle over Σ . The associated flat Lie algebra bundle will be denoted by $\text{ad } P$.

We define the parabolic cohomology, $H_{\text{par}}^1(\Sigma, \text{ad } P)$ to be the subspace of the de Rham cohomology classes in $H_{\text{DR}}^1(\Sigma, \text{ad } P)$ whose restrictions to each U_i are trivial.

7.2 Group Cohomology Construction of the Symplectic Form

Let b be the nondegenerate, G -invariant, symmetric, bilinear form on \mathfrak{g} . A skew symmetric bilinear form

$$B: H_{\text{par}}^1(\Sigma, \text{ad } P) \times H_{\text{par}}^1(\Sigma, \text{ad } P) \rightarrow H^2(\Sigma, U; \mathbb{R})$$

is defined by taking the wedge product together with the bilinear form b . Evaluating on the relative fundamental class of Σ gives the skew symmetric form,

$$A: H_{\text{par}}^1(\Sigma, \text{ad } P) \times H_{\text{par}}^1(\Sigma, \text{ad } P) \rightarrow \mathbb{R}.$$

Poincaré duality gives us nondegeneracy of A , so A is a symplectic form on $\text{Hom}(\Gamma, T; G)$. We will show A corresponds to the symplectic form $\tilde{\omega}$ given in Remark 5.3.

We first pass through the group cohomology description of $H_{\text{par}}^1(\Sigma, \text{ad } P)$ to make this correspondence explicit.

We identify the universal cover of Σ , denoted $\tilde{\Sigma}$, with the hyperbolic plane, \mathbb{H}^2 . Let $p: \tilde{\Sigma} \rightarrow \Sigma$ be the covering projection. We identify the $\mathcal{A}^\bullet(\tilde{\Sigma}, p^* \text{ad } P)$ with $\mathcal{A}^\bullet(\tilde{\Sigma}, \mathfrak{g})$ by parallel translation from a point x_0 . Given $[\eta] \in H^1(\Sigma, \text{ad } P)$ choose a representing closed 1-form $\eta \in \mathcal{A}^1(\Sigma, \text{ad } P)$. Let $\tilde{\eta} = p^* \eta$. Then there is a unique function $f: \tilde{\Sigma} \rightarrow \mathfrak{g}$ satisfying:

- $f(x_0) = 0$
- $df = \tilde{\eta}$

A 1-cochain $h(\eta) \in C^1(\Gamma, \mathfrak{g})$ is defined by

$$h(\eta)(\gamma) = f(x) - \text{Ad}_{\rho(\gamma)} f(\gamma^{-1}x).$$

This induces an isomorphism from $H^1(\Sigma, \text{ad } P)$ to $H^1(\Gamma, \mathfrak{g})$. It can be seen that $[\eta] \in H_{\text{par}}^1(\Sigma, \text{ad } P)$ if and only if $h(\eta)$ restricted to Γ_i is exact for all i . That is, there exists an $x_i \in \mathfrak{g}$ such that $h(\eta)(\gamma_i^k) = x_i - \text{Ad}_{\rho(\gamma_i^k)} x_i$ for each γ_i a generator of Γ .

We construct the fundamental domain \mathcal{D} for Γ operating on \mathbb{H}^2 as in [KM1]. Choose x_0 on Σ and make cuts along geodesics from x_0 to the cusps. The resulting fundamental domain \mathcal{D} is a geodesic $2n$ -gon with vertices v_1, \dots, v_n and cusps $v_1^\infty, \dots, v_n^\infty$ ordered so that as we proceed clockwise around $\partial \mathcal{D}$ we see $v_1, v_1^\infty, \dots, v_n, v_n^\infty$. The generator γ_i fixes v_i^∞ and satisfies $\gamma_i v_{i+1} = v_i$. Let e_i be the oriented edge joining v_i to v_i^∞ and \hat{e}_i be the oriented edge joining v_i^∞ to v_{i+1} . Then $\gamma_i \hat{e}_i = -e_i$.

Let $\rho \in \text{Hom}(\Gamma, T; G)$ and $c, c' \in T_\rho(\text{Hom}(\Gamma, T; G)/G) \simeq H_{\text{par}}^1(\Gamma, \mathfrak{g})$ be tangent vectors at ρ . The corresponding elements in $H_{\text{par}}^1(\Sigma, \text{ad } P)$ are denoted α and α' . So $f: \Sigma \rightarrow \mathfrak{g}$ which satisfies $df = \tilde{\alpha}$ and $f_i(x_0) = 0$. Let $f(v_i^\infty) = x_i$. Then

$$\begin{aligned} c(\gamma_i) &= f(x) - \text{Ad}_{\rho(\gamma_i)} f(\gamma_i^{-1}x) \\ &= f(v_i^\infty) - \text{Ad}_{\rho(\gamma_i)} f(\gamma_i^{-1}v_i^\infty) \\ &= f(v_i^\infty) - \text{Ad}_{\rho(\gamma_i)} f(v_i^\infty) \\ &= x_i - \text{Ad}_{\rho(\gamma_i)} x_i. \end{aligned}$$

There is an equivalent formulas for c' , α' , and f' with $f'(v_i^\infty) = x'_i$.

Let $B_\bullet(\Gamma)$ be the bar resolution of Γ . Thus $B_k(\Gamma)$ is the free $\mathbb{Z}[\Gamma]$ -module on the symbols $[\gamma_1|\gamma_2|\cdots|\gamma_k]$ with

$$\begin{aligned} \partial[\gamma_1|\gamma_2|\cdots|\gamma_k] \\ = \gamma_1[\gamma_2|\cdots|\gamma_k] + \sum_{i=1}^{k-1} (-1)^i [\gamma_1|\cdots|\gamma_i\gamma_{i+1}|\cdots|\gamma_k] + (-1)^k [\gamma_1|\cdots|\gamma_{k-1}]. \end{aligned}$$

Let $C_k(\Gamma) = B_k(\Gamma) \otimes_{\mathbb{Z}[\Gamma]} \mathbb{Z}$ with $\mathbb{Z}[\Gamma]$ acting on \mathbb{Z} by the homomorphism ϵ defined by

$$\epsilon\left(\sum_{i=1}^m a_i \gamma_i\right) = \sum_{i=1}^m a_i.$$

Then $C_k(\Gamma)$ is the free abelian group on the symbols $(\gamma_1|\cdots|\gamma_k) = [\gamma_1|\gamma_2|\cdots|\gamma_k] \otimes 1$ with

$$\begin{aligned} \partial(\gamma_1|\gamma_2|\cdots|\gamma_k) \\ = (\gamma_2|\cdots|\gamma_k) + \sum_{i=1}^{k-1} (-1)^i (\gamma_1|\cdots|\gamma_i\gamma_{i+1}|\cdots|\gamma_k) + (-1)^k (\gamma_1|\cdots|\gamma_{k-1}). \end{aligned}$$

A relative fundamental class $F \in C_2(\Gamma)$ is defined by the property

$$\partial F = \sum_{i=1}^n (\gamma_i).$$

Let $[\Gamma, \partial\Gamma] = \sum_{i=2}^n (\gamma_1|\cdots|\gamma_{i-1}|\gamma_i) \in C_2(\Gamma)$, then

Lemma 7.2 $[\Gamma, \partial\Gamma]$ is a relative fundamental class.

Proof The proof is left to the reader.

We will now give the symplectic form A in terms of group cohomology. We denote by \cup_b the cup product of Eilenberg-MacLane cochains using the form b on the coefficients.

Proposition 7.3

$$A(\alpha, \alpha') = \sum_{i=1}^n \langle c \cup_b x'_i, (\gamma_i) \rangle - \langle c \cup_b c', [\Gamma, \partial\Gamma] \rangle$$

We will use the next Lemmas to prove Proposition 7.3.

Lemma 7.4

$$\int_{e_i} B(f, \tilde{\alpha}') + \int_{\hat{e}_i} B(f, \tilde{\alpha}') = b(c(\gamma_i), f'(v_i^\infty)) - b(c(\gamma_i), f'(v_i))$$

Proof Recall $\gamma_i \hat{e}_i = -e_i$, so that $\hat{e}_i = -\gamma_i^{-1} e_i$. We then have

$$\begin{aligned}
\int_{e_i} B(f, \tilde{\alpha}') + \int_{\hat{e}_i} B(f, \tilde{\alpha}') &= \int_{e_i} B(f, \tilde{\alpha}') + \int_{\gamma_i^{-1} e_i} B(f, \tilde{\alpha}') \\
&= \int_{e_i} B(f, \tilde{\alpha}') + \int_{e_i} (\gamma_i^{-1})^* B(f, \tilde{\alpha}') \\
&= \int_{e_i} B(f, \tilde{\alpha}') + \int_{e_i} B((\gamma_i^{-1})^* f, (\gamma_i^{-1})^* \tilde{\alpha}') \\
&= \int_{e_i} B(f, \tilde{\alpha}') + \int_{e_i} B(\text{Ad}_{\rho(\gamma_i)}(\gamma_i^{-1})^* f, \text{Ad}_{\rho(\gamma_i)}(\gamma_i^{-1})^* \tilde{\alpha}') \\
&= \int_{e_i} B(f - \text{Ad}_{\rho(\gamma_i)}(\gamma_i^{-1})^* f, \tilde{\alpha}') \\
&= \int_{e_i} B(c(\gamma_i), \tilde{\alpha}') \\
&= b(c(\gamma_i), f'(v_i^\infty)) - b(c(\gamma_i), f'(v_i)). \quad \blacksquare
\end{aligned}$$

Lemma 7.5

$$\sum_{i=1}^n b(c(\gamma_i), f'(v_i)) = \sum_{i=1}^n b(c(\gamma_i), f'(v_i^\infty)) - \sum_{i=1}^n \langle c \cup_b y_i, (\gamma_i) \rangle + \langle c \cup_b c', [\Gamma, \partial\Gamma] \rangle.$$

Proof By definition, for any $x \in \mathbb{H}^2$ and $\gamma \in \Gamma$ we have

$$c'(\gamma) = f'(x) - \text{Ad}_{\rho(\gamma)} f'(\gamma^{-1}x).$$

Let $\gamma = \gamma_i$ and $x = v_i$, then

$$c'(\gamma_i) = f'(v_i) - \text{Ad}_{\rho(\gamma_i)} f'(v_{i+1}).$$

Using $f'(v_1) = 0$, we obtain

$$\begin{aligned}
c'(\gamma_1 \cdots \gamma_i) &= f'(v_1) - \text{Ad}_{\rho(\gamma_1 \cdots \gamma_i)} f'(\gamma_i^{-1} \cdots \gamma_1^{-1} v_1) \\
&= -\text{Ad}_{\rho(\gamma_1 \cdots \gamma_i)} f'(v_{i+1}).
\end{aligned}$$

We will also need

$$\begin{aligned}
c'(\gamma_1 \cdots \gamma_i) &= c'(\gamma_1 \cdots \gamma_{i-1}) + \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{i-1})} c'(\gamma_i) \\
&= c'(\gamma_1) + \text{Ad}_{\rho(\gamma_1)} c'(\gamma_2) + \cdots + \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{i-1})} c'(\gamma_i).
\end{aligned}$$

and, since $\gamma_1 \cdots \gamma_n = 1$,

$$0 = c'(\gamma_1 \cdots \gamma_n) = c'(\gamma_1) + \text{Ad}_{\rho(\gamma_1)} c'(\gamma_2) + \cdots + \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{n-1})} c'(\gamma_n).$$

We then have,

$$\begin{aligned}
& \sum_{i=1}^n b(c(\gamma_i), f'(v_i)) \\
&= - \sum_{i=1}^n b(c(\gamma_i), \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{i-1})} c'(\gamma_1 \cdots \gamma_{i-1})) \\
&= - \sum_{i=1}^n b\left(\text{Ad}_{\rho(\gamma_1 \cdots \gamma_{i-1})} c(\gamma_i), \sum_{j=1}^{i-1} \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{j-1})} c'(\gamma_j)\right) \\
&= - \sum_{i=1}^n \sum_{j=1}^{i-1} b(\text{Ad}_{\rho(\gamma_1 \cdots \gamma_{i-1})} c(\gamma_i), \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{j-1})} c'(\gamma_j)) \\
&= - \sum_{j=1}^n \sum_{i=j+1}^n b(\text{Ad}_{\rho(\gamma_1 \cdots \gamma_{i-1})} c(\gamma_i), \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{j-1})} c'(\gamma_j)) \\
&= \sum_{j=1}^n \sum_{i=1}^j b(\text{Ad}_{\rho(\gamma_1 \cdots \gamma_{i-1})} c(\gamma_i), \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{j-1})} c'(\gamma_j)) \\
&= \sum_{j=1}^n b(c(\gamma_1 \cdots \gamma_j), \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{j-1})} c'(\gamma_j)) \\
&= \sum_{j=1}^n b(c(\gamma_1 \cdots \gamma_{j-1}) + \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{j-1})} c(\gamma_j), \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{j-1})} c'(\gamma_j)) \\
&= \sum_{j=1}^n b(c(\gamma_1 \cdots \gamma_{j-1}), \text{Ad}_{\rho(\gamma_1 \cdots \gamma_{j-1})} c'(\gamma_j)) + \sum_{j=1}^n b(c(\gamma_j), c'(\gamma_j)) \\
&= \langle c \cup_b c', [\Gamma, \partial\Gamma] \rangle + \sum_{j=1}^n b(c(\gamma_j), f'(v_j^\infty) - \text{Ad}_{\rho(\gamma_j)} f'(v_j^\infty)) \\
&= \langle c \cup_b c', [\Gamma, \partial\Gamma] \rangle + \sum_{j=1}^n b(c(\gamma_j), f'(v_j^\infty)) - \sum_{j=1}^n \langle B(c, y_j), (\gamma_j) \rangle \quad \blacksquare
\end{aligned}$$

Proof of Proposition 7.3

$$\begin{aligned}
A(\alpha, \alpha') &= \int_{\Sigma} B(\alpha, \alpha') \\
&= \int_{\mathcal{D}} B(\tilde{\alpha}, \tilde{\alpha}') \\
&= \int_{\partial\mathcal{D}} B(\tilde{\alpha}, f')
\end{aligned}$$

$$\begin{aligned}
&= \sum_{i=1}^n \left(\int_{\hat{e}_i} B(\bar{\alpha}, f') + \int_{\hat{e}_i} B(\bar{\alpha}, f') \right) \\
&= \sum_{j=1}^n \langle c \cup_b x'_j, (\gamma_j) \rangle - \langle c \cup_b c', [\Gamma, \partial\Gamma] \rangle. \quad \blacksquare
\end{aligned}$$

7.3 Relating $\text{Hom}(\Gamma, T; \text{SU}(2)) / \text{SU}(2)$ and $M_r(\mathbb{S}^3)$

We now restrict to the case $G = \text{SU}(2)$. We define the isomorphism

$$\Upsilon: \text{Hom}(\Gamma, T; \text{SU}(2)) \rightarrow \text{CN}_r(\mathbb{S}^3, *),$$

where $\text{CN}_r(\mathbb{S}^3, *)$ is the closed polygonal linkages in \mathbb{S}^3 based at a point, by

$$\Upsilon(\rho) = (\rho(\gamma_1), \dots, \rho(\gamma_n)).$$

This induces an isomorphism, which we also denote by Υ ,

$$\Upsilon: \text{Hom}(\Gamma, T; \text{SU}(2)) / \text{SU}(2) \rightarrow M_r(\mathbb{S}^3).$$

The differential $d\Upsilon_\rho: T_\rho(\text{Hom}(\Gamma, T; \text{SU}(2)) / \text{SU}(2)) \rightarrow T_{\Upsilon(\rho)}M_r(\mathbb{S}^3)$ is then defined by

$$d\Upsilon_\rho(c) = (dR_{\rho(\gamma_1)}c(\gamma_1), \dots, dR_{\rho(\gamma_n)}c(\gamma_n)).$$

Here $T_\rho(\text{Hom}(\Gamma, T; \text{SU}(2)) / \text{SU}(2))$ is identified with an element of $\mathbb{Z}_{\text{par}}^1(\Gamma, \mathfrak{su}_2)$. We have

$$d\Upsilon_\rho(c) = (dR_{k_1}x_1 - dL_{k_1}x_1, \dots, dR_{k_n}x_n - dL_{k_n}x_n)$$

and

$$d\Upsilon_\rho(c') = (dR_{k_1}x'_1 - dL_{k_1}x'_1, \dots, dR_{k_n}x'_n - dL_{k_n}x'_n).$$

Recall, the symplectic form on $M_r(\mathbb{S}^3)$ is given by

$$\tilde{\omega} = \sum_{i=1}^n \omega_i + \frac{1}{2} \sum_{i=1}^n \sum_{j=i+1}^n (\text{Ad}_{k_1 \dots k_{i-1}} \bar{\theta}_i \wedge_b \text{Ad}_{k_1 \dots k_{j-1}} \bar{\theta}_j).$$

We can now prove the main result of this section.

Theorem 7.6 $\Upsilon^* \tilde{\omega} = A$

Proof First we note that

$$\Upsilon^* \bar{\theta}_i(c) = c(\gamma_i)$$

and

$$\begin{aligned}
(\Upsilon^* \omega_i)(c, c') &= \omega_i(dR_{k_i} c(\gamma_i), dR_{k_i} c'(\gamma_i)) \\
&= -\frac{1}{2}(\text{Ad}_{k_i^{-1}} c(\gamma_i) + c(\gamma_i), x'_i) \\
&= -\frac{1}{2}(c(\gamma_i), \text{Ad}_{k_i} x'_i + x'_i) \\
&= -\frac{1}{2}(c(\gamma_i), c'(\gamma_i)) - (c(\gamma_i), \text{Ad}_{k_i} x'_i) \\
&= -\frac{1}{2}(\text{Ad}_{k_1 \cdots k_{i-1}} c(\gamma_i), \text{Ad}_{k_1 \cdots k_{i-1}} c'(\gamma_i)) + \langle c \cup_b x'_i, (\gamma_i) \rangle.
\end{aligned}$$

It follows that

$$\begin{aligned}
(\Upsilon^* \bar{\omega})(c, c') &= \sum_{i=1}^n (\Upsilon^* \omega_i)(c, c') \\
&\quad + \frac{1}{2} \sum_{i=1}^n \sum_{j=i+1}^n \Upsilon^*(\text{Ad}_{k_1 \cdots k_{i-1}} \bar{\theta}_i \wedge_b \text{Ad}_{k_1 \cdots k_{j-1}} \bar{\theta}_j)(c, c') \\
&= \sum_{i=1}^n \langle c \cup_b x'_i, (\gamma_i) \rangle - \sum_{i=1}^n \frac{1}{2} (\text{Ad}_{k_1 \cdots k_{i-1}} c(\gamma_i), \text{Ad}_{k_1 \cdots k_{i-1}} c'(\gamma_i)) \\
&\quad + \sum_{i=1}^n \sum_{j=i+1}^n (\text{Ad}_{k_1 \cdots k_{i-1}} c(\gamma_i), \text{Ad}_{k_1 \cdots k_{j-1}} c'(\gamma_j)) \\
&\quad - \sum_{i=1}^n \sum_{j=i+1}^n (\text{Ad}_{k_1 \cdots k_{i-1}} c'(\gamma_i), \text{Ad}_{k_1 \cdots k_{j-1}} c(\gamma_j)) \\
&= \sum_{i=1}^n \langle c \cup_b x'_i, (\gamma_i) \rangle - \sum_{i=1}^n \frac{1}{2} (\text{Ad}_{k_1 \cdots k_{i-1}} c(\gamma_i), \text{Ad}_{k_1 \cdots k_{i-1}} c'(\gamma_i)) \\
&\quad + \sum_{j=2}^n \sum_{i=1}^{j-1} (\text{Ad}_{k_1 \cdots k_{i-1}} c(\gamma_i), \text{Ad}_{k_1 \cdots k_{j-1}} c'(\gamma_j)) \\
&\quad + \sum_{i=1}^n \sum_{j=1}^i (\text{Ad}_{k_1 \cdots k_{i-1}} c'(\gamma_i), \text{Ad}_{k_1 \cdots k_{j-1}} c(\gamma_j)) \\
&= \sum_{i=1}^n \langle c \cup_b x'_i, (\gamma_i) \rangle + \sum_{j=2}^n \sum_{i=1}^{j-1} (\text{Ad}_{k_1 \cdots k_{i-1}} c(\gamma_i), \text{Ad}_{k_1 \cdots k_{j-1}} c'(\gamma_j))
\end{aligned}$$

$$\begin{aligned}
&= \sum_{i=1}^n \langle c \cup_b x'_i, (\gamma_i) \rangle + \sum_{j=2}^n \langle \text{Ad}_{k_1 \dots k_{i-1}} c'(\gamma_i), c(\gamma_1 \dots \gamma_{i-1}) \rangle \\
&= \sum_{i=1}^n \langle c \cup_b x'_i, (\gamma_i) \rangle - \langle c \cup_b c', [\Gamma, \partial\Gamma] \rangle \\
&= A(\alpha, \alpha'). \quad \blacksquare
\end{aligned}$$

It is easily seen that the functions ℓ_i from Section 5.3 corresponds to the following Goldman functions. Let $\phi: \text{SU}(2) \rightarrow \mathbb{R}$ be defined by $\phi(g) = \cos^{-1}(-\frac{1}{2} \text{tr}(g))$. We then defined the function $\phi_\gamma: \text{Hom}(\Gamma, T; \text{SU}(2)) / \text{SU}(2) \rightarrow \mathbb{R}$ by $\phi_\gamma(\rho) = \phi(\rho(\gamma))$. We see that

$$\Upsilon^* \ell_i = \phi_{\gamma_1 \dots \gamma_i}.$$

Then choosing an maximal collection of nonintersecting diagonal on $M_r(\mathbb{S}^3)$ corresponds to a pair of pants decomposition on Σ .

7.4 Symplectomorphism of $M_r(\mathbb{S}^3)$ and $M_r(\mathbb{E}^3)$

We now use the following result due to L. Jeffrey.

Lemma 7.7 *There exists an open neighborhood U of 0 in \mathfrak{g}^n such that if $\bar{\lambda} = (\lambda_1, \dots, \lambda_n) \in U$ then the moduli space of parabolic bundles on n -punctured surface with weights $\lambda_1, \dots, \lambda_n$ is symplectomorphic to the symplectic reduced space $\{(X_1, \dots, X_n) \in \mathcal{O}_{\lambda_1} \times \dots \times \mathcal{O}_{\lambda_n} : X_1 + \dots + X_n = 0\} / G$.*

Proof See [Je, Theorem 6.6]. ■

We can identify the moduli space of parabolic bundles on n -punctured surface with weights $\lambda_1, \dots, \lambda_n$ with $M_r(\mathbb{S}^3)$. Also, it was shown in [KM1] that $\{(X_1, \dots, X_n) \in \mathcal{O}_{\lambda_1} \times \dots \times \mathcal{O}_{\lambda_n} \subset \mathfrak{g}^n : X_1 + \dots + X_n = 0\} / G$ can be identified with $M_r(\mathbb{E}^3)$. We have the following corollary.

Corollary 7.8 *For sufficiently small side-lengths, $M_r(\mathbb{S}^3)$ is symplectomorphic to $M_r(\mathbb{E}^3)$.*

References

- [A] A. Alekseev, *On Poisson actions of compact Lie groups on symplectic manifolds*. J. Differential Geom. **45**(1997), 241–256.
- [AKS] A. Alekseev and Y. Kosmann-Schwarzbach, *Manin pairs and moment maps*. preprint, math.DG/9909176.
- [AKSM] A. Alekseev, Y. Kosmann-Schwarzbach and E. Meinrenken, *Quasi-Poisson Manifolds*. Canad. J. Math., this issue, 3–29.
- [AMM1] A. Alekseev, A. Malkin and E. Meinrenken, *Lie group valued moment maps*. J. Differential Geom. **48**(1998), 445–495.
- [AMM2] ———, *Manin pairs of a compact simple Lie algebra*. unpublished notes.
- [Bi] J. Birman, *Braids, links, and mapping class groups*. Ann. of Math. Studies **82**, Princeton Univ. Press, 1974.

- [CP] V. Chari and A. Pressley, *A Guide to Quantum Groups*. Cambridge Univ. Press, 1994.
- [FM] H. Flaschka and J. J. Millson, *On the moduli space of n points in $\mathbb{C}P^m$* . in preparation.
- [Ga] A. Galitzer, *The moduli space of polygon linkages in the 2-sphere*. Ph.D. thesis, University of Maryland, 1997.
- [Go] W. Goldman, *Invariant functions on Lie groups and Hamiltonian flows of surface group representations*. Invent. Math. **85**(1986), 263–302.
- [GHJW] K. Guruprasad, J. Huebshmann, L. Jeffrey and A. Weinstein, *Group systems, groupoids, and moduli spaces of parabolic bundles*. Duke Math. J. **89**(1997), 377–412.
- [HK] J.-C. Hausmann and A. Knutson, *Polygon spaces and Grassmannians*. Enseign. Math. (2) **43**(1997), 173–198.
- [Je] ———, *Extended moduli spaces of flat connections on Riemann surfaces*. Math. Ann. **298**(1994), 667–692.
- [JW] L. Jeffrey and J. Weitsman, *Torus actions, moment maps, and the moduli space of flat connections on a two-manifold*. Contemp. Math. **175**(1994), 49–59.
- [KM1] M. Kapovich and J. J. Millson, *The symplectic geometry of polygons in Euclidean space*. J. Differential Geom. **44**(1996), 479–513.
- [KM2] ———, *The relative deformation theory of representations of flat connections and deformations of linkages in constant curvature spaces*. Compositio Math. **103**(1996), 287–317.
- [KM3] ———, *On the moduli space of a spherical polygonal linkage*. Canad. Math. Bull. **42**(1999), 307–320.
- [KMT] M. Kapovich, J. J. Millson and T. Treloar, *The symplectic geometry of polygons in hyperbolic 3-space*. Asian J. Math **4**(2000), 123–164.
- [KS2] Y. Kosmann-Schwarzbach, *Jacobian quasi-bialgebras and quasi-Poisson Lie groups*. Contemp. Math. **132**(1991), 459–489.
- [LM] B. Leeb and J. J. Millson, *Convex functions on symmetric spaces and geometric invariant theory for weighted configurations in flag manifolds*. in preparation.
- [Le] M. Leingang, *Symmetric pairs and moment spaces*, math.SG/9810064, preprint.
- [Lu3] J.-H. Lu, *Momentum mappings and reduction of Poisson actions*. In: Symplectic Geometry, Groupoids, and Integable Systems, MSRI Publ. **20**, Springer-Verlag, New York, 1991, 209–226.
- [Mi] J. J. Millson, *Bending polygons and decomposing tensor products, three examples*. in preparation.
- [Mo] J. Moser, *On the volume elements on a manifold*. Trans. Amer. Math. Soc. **120**(1965), 286–294.
- [MZ] J. J. Millson and B. Zombro, *A Kähler structure on the moduli space of isometric maps of a circle into Euclidean space*. Invent. Math. **123**(1996), 35–59.
- [Sa] M. Sargent, *Diffeomorphism equivalence of configuration spaces of polygons in constant curvature spaces*. Ph.D. thesis, University of Maryland, 1995.
- [STS] M. Semenov-Tian-Shansky, *Dressing transformations and Poisson group actions*. Publ. Res. Inst. Math. Sci. **21**(1985), 1237–1260.
- [Th] W. Thurston, *Three-dimensional geometry and topology*. Princeton Univ. Press, 1997.
- [Tr1] T. Treloar, *Integrable hamiltonian systems on moduli spaces of polygonal linkages*. Ph.D. thesis, University of Maryland, 2001.
- [Tr2] ———, *The symplectic geometry on loops in the 3-sphere*. in preparation.

Department of Mathematics
University of Arizona
Tucson, Arizona 85721
U.S.A.
email: treloar@math.arizona.edu