

A Note on Covering by Convex Bodies

Dedicated to Ted Bisztriczky, on his sixtieth birthday.

Gábor Fejes Tóth

Abstract. A classical theorem of Rogers states that for any convex body K in n -dimensional Euclidean space there exists a covering of the space by translates of K with density not exceeding $n \log n + n \log \log n + 5n$. Rogers' theorem does not say anything about the structure of such a covering. We show that for sufficiently large values of n the same bound can be attained by a covering which is the union of $O(\log n)$ translates of a lattice arrangement of K .

Rényi Institute of Mathematics, Hungarian Academy of Sciences, Pf. 127, H-1364 Budapest, Hungary
e-mail: gfejes@renyi.hu

Received by the editors November 12, 2007; revised April 13, 2008.

Part of this work was done during a visit of the author at the Work Group of Theoretical Computer Science in the Institute of Computer Science, Freie Universität Berlin. The visit was supported by the Discrete and Convex Geometry project (MTKD-CT-2005-014333) carried out by the Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences, in the framework of the European Community's Structuring the European Research Area programme. The research was also supported by OTKA grants T43520 and K68398.

AMS subject classification: Primary: 52C07; secondary: 52C17, 11H31.

©Canadian Mathematical Society 2009.