

THE ACADEMY CORNER

No. 38

Bruce Shawyer

All communications about this column should be sent to Bruce Shawyer, Department of Mathematics and Statistics, Memorial University of Newfoundland, St. John's, Newfoundland, Canada. A1C 5S7

In this issue, we present some readers' solutions to problems from the Memorial University Undergraduate Mathematics Competition, held in March 2000. [2000 : 257]

2. Evaluate $x^3 + y^3$ where $x + y = 1$ and $x^2 + y^2 = 2$.

Solution by Catherine Shevlin, Wallsend upon Tyne, England.

Note that

$$\begin{aligned} (x^2 + y^2)(x + y) &= x^3 + y^3 + xy(x + y) \\ &= x^3 + y^3 + \frac{1}{2}((x + y)^2 - (x^2 + y^2))(x + y). \end{aligned}$$

Therefore, $x^3 + y^3 = \frac{5}{2}$.

3. In triangle ABC , we have $\angle ABC = \angle ACB = 80^\circ$. P is chosen on line segment AB such that $\angle BPC = 30^\circ$. Prove that $AP = BC$.

Solution by D.J. Smeenk, Zaltbommel, the Netherlands.

Draw equilateral $\triangle ABQ$ with Q on the same side of AB as C .


Then $\triangle ACQ$ is isosceles, giving $AC = AQ$.

Since $\angle CAQ = 40^\circ$, we have $\angle ACQ = \angle AQC = 70^\circ$.

Similarly, since $\angle ABQ = 60^\circ$, we have $\angle CBQ = 20^\circ = \angle BAC$, so that $\angle ACP = \angle BPC - \angle BAC = 10^\circ$.

Thus, we have

$$\left. \begin{aligned} \angle CBQ &= \angle PAC = 20^\circ, \\ \angle CQB &= \angle ACP = 10^\circ, \\ BQ &= AC, \end{aligned} \right\}$$



showing that $\triangle BCQ$ is congruent to $\triangle APC$. Thus, $BC = AP$.

4. Show that $\binom{2000}{3} = (1)(1998) + (2)(1997) + \dots + k(1999 - k) + \dots + (1997)(2) + (1998)(1)$.

Solution by D.J. Smeenk, Zaltbommel, the Netherlands.

$$\begin{aligned} \sum_{k=1}^{1998} k(1999 - k) &= 2 \sum_{k=1}^{999} k(1999 - k) \\ &= 2 \cdot 1999 \cdot \sum_{k=1}^{999} k - 2 \sum_{k=1}^{999} k^2. \end{aligned} \quad (1)$$

Also,
$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}. \quad (2)$$

From (1) and (2), we obtain

$$\begin{aligned} \sum_{k=1}^{1998} k(1999 - k) &= 2 \cdot 1999 \cdot \frac{1}{2} \cdot 999 \cdot 1000 - \frac{1}{3} \cdot 999 \cdot 1000 \cdot 1999 \\ &= \frac{2}{3} \cdot 999 \cdot 1000 \cdot 1999 = \frac{1}{6} \cdot 2000 \cdot 1998 \cdot 1999 \\ &= \binom{2000}{3}. \end{aligned}$$

7. Let $f(x) = x(x-1)(x-2)\dots(x-n)$.

1. Show that $f'(0) = (-1)^n n!$
2. More generally, show that if $0 \leq k \leq n$, then $f'(k) = (-1)^{n-k} k!(n-k)!$

Solution by Catherine Shevlin, Wallsend upon Tyne, England.

Taking logarithms and differentiating, we have

$$\begin{aligned} f'(x) &= f(x) \sum_{k=0}^n \frac{1}{x-k} \\ &= (x-1)(x-2)\dots(x-n) \left(1 + \frac{x}{x-1} + \dots + \frac{x}{x-n} \right). \end{aligned}$$

Thus, $f'(0) = (-1)(-2)\dots(-n) = (-1)^n n!$.

More generally, if $0 \leq k \leq n$, we have

$$\begin{aligned} f'(x) &= x(x-1)\dots(x-k+1)(x-k-1)\dots(x-n) \\ &\quad \times \left(\frac{x-k}{x} + \dots + \frac{x-k}{x-k+1} + 1 + \frac{x-k}{x-k-1} + \dots + \frac{x-k}{x-n} \right), \end{aligned}$$

showing that

$$f'(k) = (k \cdot (k-1) \cdot \dots \cdot 2 \cdot 1) \cdot (-1) \cdot (-2) \cdot \dots \cdot (-(n-k)) = k!(-1)^{n-k}(n-k)!.$$

We end this issue with the problems of the 2000 William Lowell Putnam Mathematical Competition, reprinted with permission of the Mathematical Association of America.

**SIXTY-FIRST ANNUAL
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION
Saturday, December 2, 2000**

Problem A1.

Let A be a positive real number. What are the possible values of $\sum_{j=0}^{\infty} x_j^2$, given that x_0, x_1, x_2, \dots are positive numbers for which $\sum_{j=0}^{\infty} x_j = A$?

Problem A2.

Prove that there exist infinitely many integers n such that n , $n + 1$, and $n + 2$ are each the sum of two squares of integers.

[Example: $0 = 0^2 + 0^2$, $1 = 0^2 + 1^2$, and $2 = 1^2 + 1^2$.]

Problem A3.

The octagon $P_1P_2P_3P_4P_5P_6P_7P_8$ is inscribed in a circle, with the vertices around the circumference in the given order. Given that the polygon $P_1P_3P_5P_7$ is a square of area 5 and the polygon $P_2P_4P_6P_8$ is a rectangle of area 4, find the maximum possible area of the octagon.

Problem A4.

Show that the improper integral

$$\lim_{B \rightarrow \infty} \int_0^B \sin(x) \sin(x^2) dx$$

converges.

Problem A5.

Three distinct points with integer coordinates lie in the plane on a circle of radius $r > 0$. Show that two of these points are separated by a distance of at least $r^{1/3}$.

Problem A6.

Let $f(x)$ be a polynomial with integer coefficients. Define a sequence a_0, a_1, \dots of integers such that $a_0 = 0$ and $a_{n+1} = f(a_n)$ for all $n \geq 0$. Prove that if there exists a positive integer m for which $a_m = 0$ then either $a_1 = 0$ or $a_2 = 0$.

Problem B1.

Let $a_j, b_j,$ and c_j be integers for $1 \leq j \leq N$. Assume, for each j , that at least one of a_j, b_j, c_j is odd. Show that there exist integers r, s, t such that $ra_j + sb_j + tc_j$ is odd for at least $4N/7$ values of $j, 1 \leq j \leq N$.

Problem B2.

Prove that the expression

$$\frac{\gcd(m, n)}{n} \binom{n}{m}$$

is an integer for all pairs of integers $n \geq m \geq 1$. [Here, $\binom{n}{m} = \frac{n!}{m!(n-m)!}$ and $\gcd(m, n)$ is the greatest common divisor of m and n .]

Problem B3.

Let $f(t) = \sum_{j=1}^N a_j \sin(2\pi jt)$, where each a_j is real and $a_N \neq 0$. Let N_k

denote the number of zeros (including multiplicities) of $\frac{d^k f}{dt^k}$.

— Prove that $N_0 \leq N_1 \leq N_2 \leq \dots$ and $\lim_{k \rightarrow \infty} N_k = 2N$. —

Problem B4.

Let $f(x)$ be a continuous function such that $f(2x^2 - 1) = 2xf(x)$ for all x . Show that $f(x) = 0$ for $-1 \leq x \leq 1$.

Problem B5.

Let S_0 be a finite set of positive integers. We define finite sets S_1, S_2, \dots of positive integers as follows:

Integer a is in S_{n+1} if and only if exactly one of $a-1$ or a is in S_n .

Show that there exist infinitely many integers N for which

$$S_N = S_0 \cup \{N + a : a \in S_0\}.$$

Problem B6.

Let B be a set of more than $2^{n+1}/n$ distinct points with coordinates of the form $(\pm 1, \pm 1, \dots, \pm 1)$ in n -dimensional space, with $n \geq 3$. Show that there are three distinct points in B which are the vertices of an equilateral triangle.

Send us your interesting solutions.

We are always pleased to receive submissions for this corner.